



## Nasopharyngeal Cutaneous Tuberculosis

Polychroni D<sup>1,\*</sup> and Stefanidis I<sup>2</sup>

*Surgical Department, General Hospital of Mytilene, Lesvos, Greece*

\*Corresponding author: Polychroni D, Surgical Department, General Hospital of Mytilene, Lesvos, Greece; Email: [damaskinipi@gmail.com](mailto:damaskinipi@gmail.com)

### Clinical Image

This is a case of a 47-year-old Syrian man with a 2-year history of deep and painless nasal ulceration, causing minimal symptoms. The patient, who left his country seeking asylum and protection from human rights violation, failed to seek medical attention till he arrived to Greece. The man presented to our department with extensive and irregular ulcerative lesions, exposing his nasal septum, accompanied with pus discharge. Histopathology examination of the biopsied lesion showed Langhans-type multinucleated giant cells, surrounded by lymphocytes, revealing tuberculosis infection. The patient referred to a tertiary health care hospital to receive targeted medical therapy.



*Figure: Extensive nasal ulcerative lesions accompanied with pus discharge.*

**Received date:** 08 November 2022; **Accepted date:** 12 November 2022; **Published date:** 16 November 2022

**Citation:** Polychroni D, Stefanidis I (2022) Nasopharyngeal Cutaneous Tuberculosis. SunText Rev Case Rep Image 3(4): 161.

**DOI:** <https://doi.org/10.51737/2766-4589.2022.061>

**Copyright:** © 2022 Polychroni D, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.